

P07 43453 (18810-80645)

In the Claims:

Please amend Claims 1, 4, 9, 11-18, 21, 26, 28-34, 97, 100, 104, 108, and 109, as follows.

1. (Twice Amended) A method of delivering a medicant to an abnormal brain region in a mammalian subject, comprising:

E5 administering to a mammalian subject having an abnormal brain region an agonist of a calcium-activated or ATP-sensitive potassium channel, the agonist being other than bradykinin or a bradykinin analog, under conditions and in an amount sufficient to increase the permeability to the medicant of a capillary or arteriole delivering blood to cells of the abnormal brain region; and

administering to the subject simultaneously or substantially simultaneously with the agonist the medicant, so that the medicant is delivered selectively to the cells of the abnormal brain region compared to normal brain regions.

E6 4.(Amended) The method of Claim 1, wherein the agonist is NS-1619, 1-EBIO, a guanylyl cyclase activator, a guanylyl cyclase activating protein, minoxidil, pinacidil, cromakalim, or levcromakalim.

E7 9.(Amended) The method of Claim 1, wherein the medicant is a N-methyl-D-aspartate (NMDA) receptor antagonist, antibiotic, interleukin-2, transforming growth factor- β , cisplatin, carboplatin, tumor necrosis factor- α , methotrexate, 5-fluorouracil, amphotericin, daunorubicin, doxorubicin, vincristine, vinblastine, busulfan, chlorambucil, cyclophosphamide, melphalan, or ethyl ethanesulfonic acid.

E8 ¹⁰11.(Amended) The method of Claim 1, wherein administering the agonist is by intravenous or intra-arterial infusion or injection.

P07 43453 (18810-80645)

ES
out
11 ~~12~~. (Amended) The method of Claim 1, wherein administering the agonist is by intracarotid infusion or injection.

12 ~~13~~. (Amended) The method of Claim 1, wherein the agonist is administered to the mammalian subject by a bolus injection.

13 ~~14~~. (Amended) The method of Claim 1, wherein the agonist is administered to the mammalian subject in an amount from about 0.075 to 1500 micrograms per kilogram body mass.

14 ~~15~~. (Amended) The method of Claim ¹³~~14~~, wherein the agonist is administered to the subject in an amount from about 0.075 to 150 micrograms per kilogram body mass.

15 ~~16~~. (Amended) The method of Claim 1, wherein the agonist is administered to the mammalian subject at a dose rate of about 0.075 to about 100 $\mu\text{g kg}^{-1} \text{min}^{-1}$ for up to about 30 minutes.

16 ~~17~~. (Amended) The method of Claim ¹⁵~~16~~, wherein the agonist is administered to the mammalian subject at a dose rate of about 0.075 to about 15 $\mu\text{g kg}^{-1} \text{min}^{-1}$.

17 ~~18~~. (Twice Amended) A method of selectively delivering a medicant to an abnormal brain region in a mammalian subject, comprising:

administering to a mammalian subject having an abnormal brain region an agonist of a calcium-activated or ATP-sensitive potassium channel, the agonist being other than bradykinin or a bradykinin analog, under conditions and in an amount sufficient to increase potassium flux through a calcium-activated or ATP-sensitive potassium channel in an endothelial cell membrane of a capillary or arteriole delivering blood to cells of the abnormal brain region, whereby the capillary or arteriole is made more permeable to the medicant; and

33

Q

P07 43453 (18810-80645)

E8
Could
administering to the subject simultaneously or substantially simultaneously with the agonist the medicant, so that the medicant is delivered selectively to the cells of the abnormal brain region compared to normal brain regions.

E9
²⁰~~21~~. (Amended) The method of Claim ¹⁷~~18~~, wherein the agonist is NS-1619, 1-EBIO, a guanylyl cyclase activator, a guanylyl cyclase activating protein, minoxidil, pinacidil, cromakalim, or levcromakalim.

E10
²⁵~~26~~. (Amended) The method of Claim ¹⁷~~18~~, wherein the medicant is a N-methyl-D-aspartate (NMDA) receptor antagonist, antibiotic, interleukin-2, transforming growth factor- β , cisplatin, carboplatin, tumor necrosis factor- α , methotrexate, 5-fluorouracil, amphotericin, daunorubicin, doxorubicin, vincristine, vinblastine, busulfan, chlorambucil, cyclophosphamide, melphalan, or ethyl ethanesulfonic acid.

E11
²⁶~~28~~. (Amended) The method of Claim ¹⁷~~18~~, wherein administering the agonist is by intravenous or intra-arterial infusion or injection.

²⁷~~29~~. (Amended) The method of Claim ¹⁷~~18~~, wherein administering the agonist is by intracarotid infusion or injection.

²⁸~~30~~. (Amended) The method of Claim ¹⁷~~18~~, wherein the agonist is administered to the mammalian subject by a bolus injection.

²⁹~~31~~. (Amended) The method of Claim ¹⁷~~18~~, wherein the agonist is administered to the mammalian subject in an amount from about 0.075 to 1500 micrograms per kilogram body mass.

³⁰~~32~~. (Amended) The method of Claim ²⁹~~31~~, wherein the agonist is administered to the subject in an amount from about 0.075 to 150 micrograms per kilogram body mass.

33

Q

P07 43453 (18810-80645)

E11
Conced
31 ~~33~~ (Amended)

The method of Claim ~~18~~¹⁷, wherein the agonist is administered to the mammalian subject at a dose rate of about 0.075 to about 100 $\mu\text{g kg}^{-1} \text{min}^{-1}$ for up to about 30 minutes.

32
~~34~~ (Amended)

The method of Claim ~~33~~³¹, wherein the agonist is administered to the mammalian subject at a dose rate of about 0.075 to about 15 $\mu\text{g kg}^{-1} \text{min}^{-1}$.

F12
33
~~31~~ (Twice Amended)

A pharmaceutical composition comprising a combination of an agonist of a calcium-activated or ATP-sensitive potassium channel, the agonist being other than bradykinin or a bradykinin analog, formulated in a pharmaceutically acceptable solution together with a medicant for delivery by intravascular infusion or injection into a mammal.

F13
36 ~~100~~ (Amended)

The pharmaceutical composition of Claim ~~97~~³³, wherein the agonist is NS-1619, 1-EBIO, a guanylyl cyclase activator, a guanylyl cyclase activating protein, minoxidil, pinacidil, cromakalim, or levocromakalim.

E14
40 ~~104~~ (Twice Amended)

The pharmaceutical composition of Claim ~~97~~³³, wherein the medicant is a N-methyl-D-aspartate (NMDA) receptor antagonist, antibiotic, interleukin-2, or transforming growth factor- β .

E15
43 ~~108~~ (Thrice Amended)

A kit for enhancing the delivery of a medicant to an abnormal brain region, comprising:

an agonist of a calcium-activated or ATP-sensitive potassium channel, potassium the agonist being other than bradykinin or a bradykinin analog; and

instructions for using the agonist for enhancing the delivery of a medicant to an abnormal brain region by increasing the permeability of a capillary or arteriole delivering blood to cells of the abnormal brain region.